

# SULPHUR

## A high load formulation of elemental sulphur

### CONTENT

Sulphur (S): 800 g/l

### FORMULATION

Headland Sulphur is a high quality suspension formulation, engineered to give outstanding coverage and rapid conversion into available sulphur. The suspension is easy to pour, disperses readily and suspends well in water. It is chemically and physically stable and is compatible with a wide range of other products.

### RATES OF USE AND TIMINGS

#### Oilseed rape:

Apply 5 - 10 l/ha in at least 200 litres of water.

Make the first application just prior to stem extension. Repeat as necessary. Crops have shown good responses to foliar applied sulphur right up to mid-flowering, even in the absence of any deficiency.

#### Cereals:

Apply 5 - 10 l/ha in at least 200 litres of water. The higher rate should be used in areas with low atmospheric sulphur deposition.

#### Grassland:

Apply 5 - 10 l/ha in at least 200 litres of water.

#### Grazing grass:

Apply during early Summer. Keep livestock out of treated areas for at least 3 weeks.

#### Conservation grass:

Apply 2 weeks before the first (or second) cut.

#### Other crops:

Many other agricultural and horticultural crops such as grape vines and other fruits also show good responses to application of sulphur.

Applications should be made during periods of active growth when there is enough foliage present to absorb the spray. Do not apply when the crop is under stress due to drought or extremes of temperature.

Note: a crop's nutrient status can only be determined accurately by tissue analysis.

Headland recommends that tissue analysis results are used whenever possible to optimise micronutrient applications.

### CAUTIONS

Do not apply Sulphur when the temperature is above 25°C or in strong sunlight.

Sulphur is generally non-phytotoxic, but some varieties of apples and pears are 'sulphur shy' and care should be exercised if spraying these varieties.

It may cause taints and processors should be consulted if the crop is to be used for canning, freezing or jam making.

### APPLICATION

**Shake the container well before opening and use.** Spray equipment should be clean. Half fill the spray tank with water, begin agitation and add the required quantity of Sulphur. Rinse the container thoroughly, add the rinsings to the spray-tank, complete the filling and apply without delay. Maintain agitation throughout. Clean sprayer thoroughly after use.

### COMPATIBILITY

Sulphur is physically and chemically compatible with a wide range of plant protection products. Consult your supplier for the latest information. Up-to-date information is available at [www.headland-ag.co.uk](http://www.headland-ag.co.uk). Always read the label of the partner product carefully before mixing with Sulphur.

### STORAGE

**Protect from frost. Store above 5°C.**

Keep in cool, well ventilated suitable store. Do not store in direct sunlight or near direct sources of heat. Keep out of reach of children. Keep away from food, drink and animal feedingstuffs.

### SAFETY PRECAUTIONS

H315: Causes skin irritation.

H319: Causes serious eye irritation.

#### Warning

P264: Wash hands thoroughly after handling.

P280: Wear protective gloves/protective clothing/eye protection/face protection.

P302+352: IF ON SKIN: Wash with plenty of soap and water.

P305+351+338: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do.

Continue rinsing.

P332+313: If skin irritation occurs: Get medical advice/attention.

P337+313: If eye irritation persists: Get medical advice/attention.

Contains 1,2-Benzisothiazolin-3-one. May produce an allergic reaction.



Made in UK

Batch No: See neck of the bottle

## Contents: 10 litres